



Rhizosphere priming effects on the decomposition of soil organic matter in C₄ and C₃ grassland soils

Shenglei Fu¹ & Weixin Cheng

Department of Environmental Studies, University of California, Santa Cruz, CA 95064, USA. ¹Corresponding author*

Received 24 April 2001. Accepted in revised form 1 November 2001

Key words: ¹³C natural abundance, grassland soils, priming effect, rhizosphere, soil-derived C, soil respiration

Abstract

Using a natural abundance ¹³C method, soil organic matter (SOM) decomposition was studied in a C₃ plant – ‘C₄ soil’ (C₃ plant grown in a soil obtained from a grassland dominated by C₄ grasses) system and a C₄ plant – ‘C₃ soil’ (C₄ plant grown in a soil taken from a pasture dominated by C₃ grasses) system. In C₃ plant – ‘C₄ soil’ system, cumulative soil-derived CO₂-C were higher in the soils planted with soybean (5499 mg pot⁻¹) and sunflower (4484 mg pot⁻¹) than that in ‘C₄ soil’ control (3237 mg pot⁻¹) without plants. In other words, the decomposition of SOM in soils planted with soybean and sunflower were 69.9% and 38.5% faster than ‘C₄ soil’ control. In C₄ plant – ‘C₃ soil’ system, there was an overall negative priming effect of live roots on the decomposition of SOM. The cumulative soil-derived CO₂-C were lower in the soils planted with sorghum (2308 mg pot⁻¹) and amaranthus (2413 mg pot⁻¹) than that in ‘C₃ soil’ control (2541 mg pot⁻¹). The decomposition of SOM in soils planted with sorghum and amaranthus were 9.2% and 5.1% slower than ‘C₃ soil’ control. Our results also showed that rhizosphere priming effects on SOM decomposition were positive at all developmental stages in C₃ plant – ‘C₄ soil’ system, but the direction of the rhizosphere priming effect changed at different developmental stages in the C₄ plant – ‘C₃ soil’ system. Implications of rhizosphere priming effects on SOM decomposition were discussed.

Introduction

The rhizosphere has been recognized as one of the key fine scale components in global carbon cycle research (Coleman et al., 1992). Many important aspects of plant–soil interactions such as plant nutrient acquisition (Uren and Reisenauer, 1988), root colonization by rhizosphere microorganisms (Baker, 1991), and soil organic matter (SOM) decomposition (Cheng and Coleman, 1990; Sallih and Bottner, 1988) are mediated by rhizosphere processes. According to Kuzyakov et al. (2000), priming effects are strong short-term changes in the turnover of soil organic matter caused by comparatively moderate treatments (or disturbance) of the soil. Planting is one of the treatments (or disturbance) of soil, and plant root exudation or rhizosphere deposition may induce rhizo-

sphere priming effect. In the literature, both positive (Helal and Sauerbech, 1984, 1987; Sallih and Bottner, 1988) and negative (Cheng, 1996; Liljieroth et al., 1994) rhizosphere priming effects on SOM decomposition have been reported. The contrasting findings may be explained by the link between SOM decomposition processes, plant carbon allocation and the metabolic status of the rhizosphere microbial community (Cheng and Coleman, 1990). For example, carbon allocation belowground for root growth and exudation may enhance microbial growth resulting in an increase in SOM decomposition. In contrast, plant uptake for nutrients might suppress soil microbial growth in nutrient-limiting soils, consequently, reduces decomposition of SOM (Hu et al., 2001).

The effects of plant roots on SOM decomposition have been studied using ¹⁴C labeling technique in two approaches: (1) Label plants with ¹⁴C in a soil which was not previously ¹⁴C labeled (Helal and

* E-mail: fushenglei@hotmail.com

Sauerback, 1984, 1987; Liljieroth et al., 1994.). (2) Apply ^{14}C labeled plant residue or other material (e.g. ^{14}C labeled glucose) into soil, and it decomposes with the presence of live plants (Cheng and Coleman, 1990; Helal and Sauerback, 1987; Jenkinson, 1977; Reid and Goss, 1982; Sallih and Bottner, 1988). Both approaches enable one to differentiate the origin of $\text{CO}_2\text{-C}$ evolved from rhizosphere or from SOM. The $^{14}\text{CO}_2$ was considered to be that from the decomposition or utilization of ^{14}C labeled plant materials (residues or root exudates) and the $^{12}\text{CO}_2$ was from the decomposition or utilization of non-labeled materials (e.g. SOM). Natural abundance ^{13}C tracer method has many advantages over radioactive ^{14}C labeling technique (Cheng, 1996). However, ^{13}C natural tracer method has not been a popular approach for decomposition studies until recent years (Cheng, 1996; Ekblad and Högberg, 2000; Nyberg et al., 2000; Robinson and Scrimgeour, 1995; Rochette and Flanagan, 1997; Rochette et al., 1999). Isotope ^{13}C techniques have been employed in decomposition studies in two ways: (1) growing C_4 plants in 'C₃ soil' (derived from C_3 plant) or vice versa (Cheng, 1996; Robinson and Scrimgeour, 1995; Rochette and Flanagan, 1997; Rochette et al., 1999). (2) and apply C_4 plant residues or other C_4 materials (e.g. C_4 -sucrose) on 'C₃ soil' or vice versa without the presence of live plants (Ekblad and Högberg, 2000; Nyberg et al., 2000). Unfortunately, these studies have not been able to estimate the rhizosphere priming effects on SOM decomposition either because of the absence of live plants (Nyberg et al., 2000) or lack of data from control soils without live plants (Rochette and Flanagan, 1997; Robinson and Scrimgeour, 1995).

In the present study, we established a C_3 plant – 'C₄ soil' and a C_4 plant – 'C₃ soil' systems and investigated the effects of plant roots on SOM decomposition at different plant developmental stage. The soil-derived C to total soil respiration was partitioned using the ^{13}C natural tracer method, and was used as a measure of the decomposition of SOM. Our objective was to test the hypothesis that rhizosphere priming effects on SOM decomposition vary with plant species and plant developmental stages in both C_3 plant – 'C₄ soil' and C_4 plant – 'C₃ soil' systems.

Materials and methods

Soils

Two types of soils were selected for this study. One soil was obtained from Konza Prairie, KS, where C_4 grasses dominated, and the soil was then defined as 'C₄ soil' in the present study. 'C₄ soil' had a $\delta^{13}\text{C}$ of -14.61‰ , C and N contents of 20 g kg^{-1} and 1.9 g kg^{-1} , and a pH of 7.05. The other soil was taken from a Santa Cruz pasture, CA, where C_3 grasses dominated, and it was defined as 'C₃ soil'. The $\delta^{13}\text{C}$ value and soil pH of 'C₃ soil' were -24.16‰ , 5.49, and soil C and N contents were 19 g kg^{-1} and 1.7 g kg^{-1} , respectively. The soils were sieved with a 4-mm mesh sieve and air-dried before use.

Treatments

PVC (polyvinyl chloride) pots (15 cm in diameter, 40 cm in height, capped bottom, with air outlet tubing at the bottom) were made and used for growing plants. Eight kg of air-dried Kansas 'C₄ soil' or Santa Cruz 'C₃ soil' was put into each PVC pot. Soil moisture of these soils was adjusted to 300 g kg^{-1} prior to planting. Two C_3 plant species, soybean (*Glycine max*) and sunflower (*Helianthus annuus*), were grown in 'C₄ soil', and two C_4 plants, sorghum (*Sorghum bicolor*) and amaranthus (*Amaranthus hypochondriacus*) were grown in 'C₃ soil'. Planting was conducted on March 27, 2000 with multiple seeds but only one plant in each pot was kept for later measurements. Controls for 'C₄ soil' or 'C₃ soil' were set in the same way except without plants. Soil water content in each pot was maintained at about 300 g kg^{-1} by weighing and watering.

This experiment was conducted in a greenhouse of the Department of Biology at University of California, Santa Cruz. During the period of plant growth, photoperiod was set as 14 h with supplement lighting when needed. Temperature was maintained at 22 °C to 25 °C , relative humidity was maintained at 450 g kg^{-1} during the day and 750 g kg^{-1} during the night.

Sampling and measurements

Sampling was conducted on May 10 and 11, May 31 and June 1, June 25 and 26, and July 18 and 19, 2000, representing the vegetative, flowering, grain-filling and maturity stages of the plants, respectively. There were four replication pots for each treatment at

each sampling time. Sampling included CO₂ trapping, plant biomass harvesting and soil collection.

For CO₂ trapping, two pieces of half-moon-shaped Plexiglas sheets connected with Tygon tubing (3.2 mm I.D.) were placed and fitted around the plant base and taped together. Low melting point paraffin wax was melted and poured on the top of the Plexiglas sheets to seal the top of the PVC pot. The Tygon tubing connected to the Plexiglas sheet served as the air inlet during CO₂ trapping. Pots were tested for possible air leaks by pumping air into the pot through the air inlet and submerging the air outlet tubing into water. The CO₂ accumulated in the PVC pot was completely pumped out before the start of the CO₂ trapping. All CO₂ evolved from the plant–soil system were trapped for two consecutive diurnal cycles (or 48 h) with a closed circulating system (Cheng, 1996). The closed circulating system consisted of three major components: Plant–soil system, CO₂ absorber, and an airflow controlling and pressure relief set. All gases (including CO₂) evolved from the plant–soil system were pumped through a CO₂ trapping column (a PVC tube filled with a mixture of 35 ml of 4 N NaOH solution and 350 g burnt and acid-washed fine sand). The coating of NaOH on fine sand particles insured a complete trapping of CO₂, no CO₂ was detected with infrared gas analyzer (IRGA) after all gases traveled through the sand column. The rest of the gases passed a needle valve, a flow meter and a Mylar balloon (filled with CO₂-free air, served as a pressure relief and oxygen source), and was brought back to the plant–soil system. Possible air contamination during the trapping process was corrected by setting the blank CO₂ traps with the same set-ups but bypassing the plant–soil system. The CO₂ trapping columns were brought back to laboratory and the content of each column was washed with deionized water and collected in a sample bottle. After settled down, the inorganic C content of a sub-sample of the solution was measured with a TOC analyzer (Shimadzu TOC-5050A). Another sub-sample was used to form the SrCO₃ precipitate with SrCl₂, excess SrCl₂ was added to insure a complete precipitation of the carbonate. The SrCO₃ precipitate was washed with deionized water until its pH was 7 to minimize the air contamination to the sample (preventing further absorption of air CO₂) and then dried in an oven (Harris, 1997).

Immediately after CO₂ trapping, plant materials were harvested and categorized into stem, leaf and root. Plant roots were washed (e.g. roots) with deionized water and collected in brown paper bags, and

dried in an oven at 70 °C. One composite soil sample was taken from each PVC pot and a sub-sample was dried in an oven at 105 °C.

All oven-dried materials (SrCO₃ precipitate, plant materials and soils) were ground with a Spex mill (Certiprep 8000), and their δ¹³C were measured with a PDZ Europa (Cheshire, UK) 'Hydra 20-20' continuous flow isotope ratio mass spectrometer. The samples were combusted at 1000 °C and the combustion products (N₂ and CO₂) fed into the IRMS via a chromatography column which separated N₂ and CO₂. The isotope ratios were compared to those of reference gas injections.

Calculation

If one grows C₃ plant in a 'C₄ soil', the carbon derived from C₃ plant (δ¹³C = -24‰ to -29‰) and from 'C₄ soil' (δ¹³C = -12‰ to -14‰) will have different δ¹³C values. Plant derived CO₂-C was estimated using the following equation:

$$C_3 = C_t (\delta_t - \delta_4) / (\delta_3 - \delta_4)$$

In the equation, C_t is the total C from soil respiration (C_t = C₃ + C₄), C₃ is the amount of C derived from C₃ plant roots, C₄ is the amount of C derived from decomposition of SOM in 'C₄ soil', δ_t is the δ¹³C value of C_t, δ₃ is the δ¹³C value of the C derived from C₃ plant roots. Cheng (1996) found that total rhizosphere CO₂ has the same δ¹³C value as that of plant roots, i.e. isotope fractionation does not occur during rhizosphere respiration. δ₄ is the δ¹³C value of the C derived from decomposition of SOM in 'C₄ soil'. Likewise, if one grows C₄ plant in a 'C₃ soil', plant derived CO₂-C can be estimated using a similar equation:

$$C_4 = C_t (\delta_t - \delta_3) / (\delta_4 - \delta_3)$$

Here, C_t is the total C from soil respiration (C_t = C₃ + C₄), C₄ is the amount of C derived from C₄ plant roots, C₃ is the amount of C derived from decomposition of SOM in 'C₃ soil', δ_t is the δ¹³C value of C_t, δ₄ is the δ¹³C value of the C derived from C₄ plant roots, and δ₃ is the δ¹³C value of the C derived from decomposition of SOM in 'C₃ soil'.

Cumulative soil-derived CO₂-C through the whole experiment was estimated by extrapolation and it was the sum of the cumulative soil-derived CO₂-C of all plant developmental stages. Cumulative soil-derived CO₂-C for a specific developmental stage was estimated by multiplying the duration (days) with the mean value of daily soil-derived CO₂-C of the stage.




Figure 1. Daily soil-derived CO₂-C under different plant-soil systems. KC – Kansas 'C₄ soil' control; KSB – Soybean planted on 'C₄ soil'; KSF – Sunflower planted on 'C₄ soil'; SC – Santa Cruz 'C₃ soil' control; SSG – Sorghum planted on 'C₃ soil'; SAR – Amaranthus planted on 'C₃ soil'. V, F, G and M represent the vegetative, flowering, grain-filling and maturity stages of the plants. The error bars are standard errors.

The difference in cumulative soil-derived CO₂-C between a soil with and without plant was defined as rhizosphere-primed soil C. For instance, the rhizosphere-primed soil C under the influence of soybean was calculated by subtracting the cumulative soil-derived CO₂-C of 'C₄ soil' control from the cumulative soil-derived CO₂-C of that soil planted with soybean.

Statistics analysis

Statistical analyses for all data were performed using SAS software (SAS Institute, 1985) GLM procedure. Comparison among means was carried out using the LSD test for equal sample sizes and Scheffe's test for unequal sample sizes. Significance levels were set at $P < 0.05$. Unless otherwise stated, all data were expressed on dry mass basis.

Results

Daily soil-derived CO₂-C under different plant-soil systems

Daily soil-derived CO₂-C was the portion of soil respiration that originated from the decomposition of SOM. In C₃ plant – 'C₄ soil' system, daily soil-derived CO₂-C were significantly higher from the soils planted with soybean and sunflower than from 'C₄ soil' control during all developmental stages. Daily soil-derived CO₂-C from the treatment of soybean was significantly lower at vegetative stage than that at later stages. There was no significant difference




Figure 2. Cumulative soil-derived CO₂-C under different plant-soil systems. KC – Kansas 'C₄ soil' control; KSB – Soybean planted on 'C₄ soil'; KSF – Sunflower planted on 'C₄ soil'; SC – Santa Cruz 'C₃ soil' control; SSG – Sorghum planted on 'C₃ soil'; SAR – Amaranthus planted on 'C₃ soil'. V, F, G and M represent the vegetative, flowering, grain-filling and maturity stages of the plants.

in daily soil-derived CO₂-C between different developmental stages for the soils planted with sunflower (Figure 1). In C₄ plant – 'C₃ soil', the daily soil-derived CO₂-C was significantly less than that of 'C₃ soil' control at grain-filling stage for sorghum and at vegetative stage for amaranthus, but significantly more at maturity stage for amaranthus (Figure 1).

Cumulative soil-derived CO₂-C and SOM decomposition

The pattern of cumulative soil-derived CO₂-C reflected the dynamics of the decomposition of SOM. In C₃ plant – 'C₄ soil' system, the cumulative soil-derived CO₂-C were higher in the soils with live soybean and sunflower roots than that in 'C₄ soil' control at all developmental stages, illustrating a positive priming effect of live roots on the decomposition of SOM. Over the whole experimental period, the cumulative soil-derived CO₂-C were 5499 and 4484 mg pot⁻¹ for soils planted with soybean and sunflower, indicating that the SOM decomposition in soils planted with soybean and sunflower were 69.9% and 38.5% faster than that in 'C₄ soil' control (3237 mg pot⁻¹) (Figure 2). Unlike in C₃ plant – 'C₄ soil' system, the direction of rhizosphere priming effect changed at different plant developmental stages in C₄ plant – 'C₃ soil' system. The cumulative soil-derived CO₂-C were lower at vegetative, grain-filling and maturity stages but higher at flowering stage in the soil with live sorghum roots than that in 'C₃ soil' control. The cumulative soil-derived CO₂-C were lower at vegetative and flowering stages but higher at grain-filling and maturity stages in the

Table 1. Correlation between root biomass and rhizosphere-primed soil C

System	R^2				P value			
	V ^a	F	G	M	V	F	G	M
C ₃ plant – ‘C ₄ soil’	0.06 ^b	0.37	0.32	0.62	0.84	0.09*	0.14	0.02**
C ₄ plant – ‘C ₃ soil’	0.02	0.09	0.58	0.61	0.79	0.47	0.03**	0.02**
C ₃ plant – ‘C ₄ soil’	0.28 (n=32)			0.0018***				
C ₄ plant – ‘C ₃ soil’	0.05 (n=32)			0.7042				

^aV, F, G and M represent vegetative, flowering, grain-filling and maturity stages of plants, respectively.

^bn = 8.

soil with live amaranthus roots than that in ‘C₃ soil’ control. Over the whole experimental period, cumulative soil-derived CO₂-C were 2308 and 2413 mg pot⁻¹ for soils planted with sorghum and amaranthus which suggested that the SOM decomposition in soils planted with sorghum and amaranthus were 9.2% and 5.1% slower than that of ‘C₃ soil’ control (2541 mg pot⁻¹)(Figure 2).

Correlation between rhizosphere-primed soil C and root biomass

In C₃ plant – ‘C₄ soil’ system, there was a significant positive correlation between rhizosphere-primed soil C and root biomass at flowering and maturity stages. In C₄ plant – ‘C₃ soil’ system, the correlation was significantly negative at grain-filling and maturity stages. Overall, there was a significantly positive correlation between rhizosphere-primed soil C and root biomass in C₃ plant – ‘C₄ soil’ system; and no significant correlation in C₄ plant – ‘C₃ soil’ system (Table 1).

Discussion

Helal and Sauerbeck (1984, 1987) reported that maize roots significantly enhanced the breakdown of SOM, and thus induced a positive priming effect. However, Liljieroth et al. (1994) found that maize did not cause priming effect on SOM, but they found that wheat suppressed the decomposition of SOM at high N level. Cheng (1996) also found that the presence of living wheat roots suppressed the decomposition of original soil C. Sallih and Bottner (1988) reported that the system with plants lost significantly more native soil organic C than the unplanted control for one soil type but this was not observed for another soil. Our results showed that the overall rhizosphere priming effects

on the decomposition of SOM were positive (38.5–69.9% more) for a ‘C₄ soil’ planted with sunflower and soybean and were negative (5.1–9.2% less) for a ‘C₃ soil’ planted with amaranthus and sorghum over the whole experimental period. Plant species (Liljieroth et al., 1994), growth conditions (Helal and Sauerbeck, 1984, 1987; Liljieroth et al., 1994), soil nutrient level (Hu, 2001; Wang and Bakken, 1997) and soil types (Kuzyakov et al., 2000; Sallih and Bottner, 1988) seemed to be the factors affecting the direction (positive or negative) of the priming effect. However, none of the studies mentioned the plant phenology. The results of cumulative soil-derived CO₂-C from the present study illustrated that the direction of priming effect of live roots on the decomposition of SOM changed with plant developmental stages in C₄ plant – ‘C₃ soil’ system (Figure 2). This might be one of the causes for contrasting findings in different studies if the measurements were made at different developmental stages of the plants. Our results suggested that not only plant species but also plant phenology should be taken into account in the studies of rhizosphere priming effects on SOM decomposition.

Rhizodeposition has been shown to be a primer for the mineralization of SOM (Helal and Sauerbeck, 1986). Rhizodeposition includes all the losses of organic substances arising from the plant roots. It consists of water-soluble root exudates, secretion, lysates (e.g. cell walls, sloughed cells and decaying roots) (Whipps, 1990; Whipps and Linch, 1985). Rhizodeposition, as easily decomposable organic substances, can increase microbial activity or microbial biomass turnover, therefore accelerate SOM mineralization (positive priming effect). However, rhizodeposition can be toxic substances to microorganisms, therefore inhibit the microbial activity or their enzymes and suppress SOM decomposition (negative priming effect). The significant correlation between rhizosphere-primed soil C and root biomass at certain plant developmental stages supports that the rhizodeposition is one of the causes for priming effect. Nevertheless, rhizosphere priming effect can not be attributed to rhizodeposition alone. Roots exploration into soil can destabilize soil aggregates and thereby releases previously protected soil organic carbon, and the growth of roots also increases dry-wet cycles and therefore affects the decomposition process of soil organic carbon (Helal and Sauerbeck, 1987). In addition, competition between living roots and rhizosphere micro-organisms for limited nutrients (i.e. nitrogen) might suppress microbial growth and cause negative priming effect

(Cheng, 1996; Hu et al., 2001; Kuzyakov et al., 2000; Wang and Bakken, 1997). The exact cause for the shift of the direction of rhizosphere priming effect at different plant developmental stages can not be sorted out without simultaneous measurements of the parameters such as: quantity and quality of rhizodeposition, soil moisture and C/N of microbial biomass.

'Litterbag' is still one of the most popular methods employed in decomposition studies (Heneghan et al., 1999; Melillo et al., 1982), and often with the absence of live plants. The major drawback of the 'litterbag' method is that it ignores the influence of plant roots on the decomposition of SOM. Our findings also illustrate that careful assumptions (i.e. if rhizosphere priming effect is negligible) must be made for the use of root exclusion method to estimate root respiration under field conditions. Our results suggest that the rhizosphere priming effect on SOM decomposition should be taken into account in further C budgeting and modeling.

Acknowledgements

We acknowledge John Blair for helping us obtain soil from Konza prairie, Kansas. We thank Rick Susfalk for his help in greenhouse and his comments on the manuscript. We also thank Rick Terry, Jakob Rowny, Michael Lebeck, Melinda Leash, Carly Boyd, Christina Dell and Peter Zeites for their greenhouse and laboratory assistance. The comments from the editor and two anonymous reviewers greatly improved the manuscript. This research was supported by a CSREES/NRI grant from the United States Department of Agriculture (Award # 98-35107-7013).

References

- Baker R 1991 Induction of rhizosphere competence in the biocontrol fungus *Trichoderma*. In *The Rhizosphere and Plant Growth*. Eds. D L Keister and P B Cregan. pp 221–228. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Cheng W 1996 Measurement of rhizosphere respiration and organic matter decomposition using natural ^{13}C . *Plant Soil* 183, 163–168.
- Cheng W and Coleman D C 1990 Effect of living roots on soil organic matter decomposition. *Soil Biol. Biochem.* 22, 781–787.
- Coleman D C, Odum E P and Crossley Jr D A 1992 Soil biology, soil ecology and global change. *Biol. Fert. Soils* 14, 104–111.
- Ekblad A and Högberg P 2000 Analysis of $\delta^{13}\text{C}$ of CO_2 distinguishes between microbial respiration of added C_4 -sucrose and other soil respiration in a C_3 -ecosystem. *Plant Soil* 219, 197–209.
- Harris D, Porter L K and Paul E A 1997 Continuous flow isotope ratio mass spectrometry of carbon dioxide trapped as strontium carbonate. *Commun. Soil Sci. Plant Anal.* 28, 747–757.
- Helal H M and Sauerback D R 1984 Influence of plant roots on C and P metabolism in soil. *Plant Soil* 76, 175–182.
- Helal H M and Sauerback D R 1987 Direct and indirect influence of plant roots on organic matter and phosphorus turnover in soil. *INTECOL Bull.* 15, 49–58.
- Heneghan L, Coleman D C, Zou X, Crossley Jr D A and Haines B L 1999 Soil microarthropod contributions to decomposition dynamics: Tropical-temperate comparisons of single substrate. *Ecology* 80, 1873–1882.
- Hu S, Chapin III F S, Firestone M K, Field C B and Chiariello N R 2001 Nitrogen limitation of microbial decomposition in a grassland under elevated CO_2 . *Nature* 409, 188–191.
- Jenkinson D S 1977 Studies on the decomposition of plant material in soil. V. The effects of plant cover and soil type on the loss of carbon from ^{14}C labelled ryegrass decomposing under field conditions. *J. Soil Sci.* 28, 424–434.
- Kuzyakov Y, Friedel J K and Stahr K 2000 Review of mechanisms and quantification of priming effects. *Soil Biol. Biochem.* 32, 1485–1498.
- Liljieroth E, Kuikman P and Van Veen J A 1994 Carbon translocation to the rhizosphere of maize and wheat and influence on the turnover of native soil organic matter at different soil nitrogen levels. *Plant Soil* 161, 233–240.
- Melillo J M, Aber J D and Muratore J F 1982 Nitrogen and lignin control of hardwood leaf litter decomposition dynamics. *Ecology* 63, 621–626.
- Nyberg, Ekblad A, Buresh R J and Högberg P 2000 Respiration from C_3 plant green manure added to a C_4 plant carbon. *Plant Soil* 218, 83–89.
- Reid J B and Goss M J 1982 Suppression of decomposition of ^{14}C -labelled plant roots in the presence of living roots of maize and perennial ryegrass. *J. Soil Sci.* 33, 387–395.
- Robinson D and Scrimgeour C M 1995 The contribution of plant C to soil CO_2 measured using $\delta^{13}\text{C}$. *Soil Biol. Biochem.* 27, 1653–1656.
- Rochette P and Flanagan L B 1997 Quantifying rhizosphere respiration in a corn crop under field conditions. *Soil Sci. Soc. Am. J.* 61, 466–474.
- Rochette P, Flanagan L B and Gregorich E G 1999 Separating soil respiration into plant and soil components using analyses of natural abundance of carbon-13. *Soil Sci. Soc. Am. J.* 63, 1207–1213.
- Sallih Z and Bottner P 1988 Effects of wheat (*Triticum aestivum*) roots on mineralization rates of soil organic matter. *Biol. Fert. Soil.* 7, 67–70.
- SAS Institute Inc. 1985. *SAS Procedures Guide for Personal Computers*, Version 6. Edition. Cary, North Carolina.
- Uren N C and Reisenauer H M 1988 The role of root exudates in nutrient acquisition. *Advances in Plant Nutrition* 3, 79–114.
- Wang J G and Bakken L R 1997 Competition for nitrogen during mineralization of plant residues in soil: microbial response to C and N availability. *Soil Biol. Biochem.* 29, 163–170.
- Whipps J M 1990 Carbon economy. In *The Rhizosphere*. Ed. J M Lynch. pp 59–97. Wiley and Sons, Chichester, UK.
- Whipps J M and Lynch J M 1985 Energy losses by the plant in rhizodeposition. *Ann. Proc. Phytochem. Soc. Eur.* 26, 59–71.

Section editor: R. Merckx