

5-1. If E is a vector bundle over a topological space M , show that the projection map $\pi: E \rightarrow M$ is a homotopy equivalence.

5-2. Prove that the space E constructed in Example 5.2, together with the projection $\pi: E \rightarrow \mathbb{S}^1$, is a smooth rank-1 vector bundle over \mathbb{S}^1 , and show that it is nontrivial.

5-3. Let $\pi: E \rightarrow M$ be a smooth vector bundle of rank k over a smooth manifold M . Suppose $\{U_\alpha\}_{\alpha \in A}$ is an open cover of M , and for each $\alpha \in A$ we are given a smooth local trivialization $\Phi_\alpha: \pi^{-1}(U_\alpha) \rightarrow U_\alpha \times \mathbb{R}^k$ of E . For each $\alpha, \beta \in A$ such that $U_\alpha \cap U_\beta \neq \emptyset$, let $\tau_{\alpha\beta}: U_\alpha \cap U_\beta \rightarrow \text{GL}(k, \mathbb{R})$ be the transition function defined by (5.3). Show that the following identity is satisfied for all $\alpha, \beta, \gamma \in A$:

$$\tau_{\alpha\beta}(p)\tau_{\beta\gamma}(p) = \tau_{\alpha\gamma}(p), \quad p \in U_\alpha \cap U_\beta \cap U_\gamma. \quad (5.6)$$

(Here juxtaposition of matrices represents matrix multiplication.)

5-4. Let M be a smooth manifold and let $\{U_\alpha\}_{\alpha \in A}$ be an open cover of M . Suppose for each $\alpha, \beta \in A$ we are given a smooth map $\tau_{\alpha\beta}: U_\alpha \cap U_\beta \rightarrow \text{GL}(k, \mathbb{R})$ such that (5.6) is satisfied for all $\alpha, \beta, \gamma \in A$. Show that there is a smooth rank- k vector bundle $E \rightarrow M$ with smooth local trivializations $\Phi_\alpha: \pi^{-1}(U_\alpha) \rightarrow U_\alpha \times \mathbb{R}^k$ whose transition functions are the given maps $\tau_{\alpha\beta}$. [Hint: Define an appropriate equivalence relation on $\coprod_{\alpha \in A} (U_\alpha \times \mathbb{R}^k)$, and use the bundle construction lemma.]

5-5. Let $\pi: E \rightarrow M$ and $\tilde{\pi}: \tilde{E} \rightarrow M$ be two smooth rank- k vector bundles over a smooth manifold M . Suppose $\{U_\alpha\}_{\alpha \in A}$ is an open cover of M such that both E and \tilde{E} admit smooth local trivializations over each U_α . Let $\{\tau_{\alpha\beta}\}$ and $\{\tilde{\tau}_{\alpha\beta}\}$ denote the transition functions determined by the given local trivializations of E and \tilde{E} , respectively. Show that E and \tilde{E} are smoothly isomorphic over M if and only if for each $\alpha \in A$ there exists a smooth map $\sigma_\alpha: U_\alpha \rightarrow \text{GL}(k, \mathbb{R})$ such that

$$\tilde{\tau}_{\alpha\beta}(p) = \sigma_\alpha(p)^{-1}\tau_{\alpha\beta}(p)\sigma_\beta(p), \quad p \in U_\alpha \cap U_\beta.$$

5-6. Let $U = \mathbb{S}^1 \setminus \{1\}$ and $V = \mathbb{S}^1 \setminus \{-1\}$, and define $\tau: U \cap V \rightarrow \text{GL}(1, \mathbb{R})$ by

$$(1) \quad \text{Im } z > 0,$$

that F is smoothly isomorphic to the Möbius bundle of Example 5-2

- 5-7. Compute the transition function for TS^2 associated with the two local trivializations determined by stereographic coordinates.
- 5-8. Let $\pi: E \rightarrow M$ be a smooth vector bundle of rank k , and suppose $\sigma_1, \dots, \sigma_m$ are independent smooth local sections over an open subset $U \subset M$. Show that for each $p \in U$ there are smooth sections $\sigma_{m+1}, \dots, \sigma_k$ defined on some neighborhood V of p such that $(\sigma_1, \dots, \sigma_k)$ is a smooth local frame for E over $U \cap V$.
- 5-9. Suppose E and E' are vector bundles over a smooth manifold M , and $F: E \rightarrow E'$ is a bijective bundle map over M . Show that F is a bundle isomorphism.
- 5-10. Consider the following vector fields on \mathbb{R}^4 :

$$\begin{aligned}X_1 &= -x^2 \frac{\partial}{\partial x^1} + x^1 \frac{\partial}{\partial x^2} + x^4 \frac{\partial}{\partial x^3} - x^3 \frac{\partial}{\partial x^4}, \\X_2 &= -x^3 \frac{\partial}{\partial x^1} - x^4 \frac{\partial}{\partial x^2} + x^1 \frac{\partial}{\partial x^3} + x^2 \frac{\partial}{\partial x^4}, \\X_3 &= -x^4 \frac{\partial}{\partial x^1} + x^3 \frac{\partial}{\partial x^2} - x^2 \frac{\partial}{\partial x^3} + x^1 \frac{\partial}{\partial x^4}.\end{aligned}$$

Show that there are smooth vector fields V_1, V_2, V_3 on S^3 such that V_j is ι -related to X_j for $j = 1, 2, 3$, where $\iota: S^3 \hookrightarrow \mathbb{R}^4$ is inclusion. Conclude that S^3 is parallelizable.

- 5-11. Let V be a finite-dimensional vector space, and let $G_k(V)$ be the Grassmannian of k -dimensional subspaces of V . Let T be the disjoint union of all these k -dimensional subspaces:

$$T = \coprod_{S \in G_k(V)} S;$$

and let $\pi: T \rightarrow G_k(V)$ be the natural map sending each point $x \in S$ to S . Show that T has a unique smooth manifold structure making it into a smooth rank- k vector bundle over $G_k(V)$, with π as projection and with the vector space structure on each fiber inherited from V . [Remark: T is sometimes called the *tautological vector bundle* over $G_k(V)$, because the fiber over each point $S \in G_k(V)$ is S itself.]

- 5-12. Show that the tautological vector bundle over $G_1(\mathbb{R}^2)$ is isomorphic to the Möbius bundle. (See Problems 5-2, 5-6, and 5-11.)
- 5-13. Let \mathcal{V}_0 be the category whose objects are finite-dimensional real vector spaces and whose morphisms are linear isomorphisms. If \mathcal{F} is a covariant functor from \mathcal{V}_0 to itself, for each finite-dimensional vector