

Manifolds 1 Homework 1

Elijah Fender

October 10, 2016

To show that N is diffeomorphic to the mbius band we define a smooth map from N to \mathbb{RP}^2 minus a point. And to do this we define a map from N to $\mathbb{R}^3 \setminus 0$ that is constant on the fibers of the quotient map from $\mathbb{R}^3 \rightarrow \mathbb{RP}^2$ which we will denote π . First note that we can write any line in N in the form $ax + by + c = 0$ as long as a and b are not both zero. Now note that $\lambda(ax + by + c = 0) = ax + by + c$. That is to say multiplying this equation by a scalar doesn't change what line it is in the plane. Now we define $\phi : N \rightarrow \mathbb{R}^3 \setminus 0$ by $ax + by + c = 0 \mapsto [a, b, c]$. As stated above this function's inverse image on linear subspaces of \mathbb{R}^3 map to a single line (which is a point in N) showing that ϕ is constant on the fibers of π . It follows that this descends to a smooth map $\pi \circ \phi : N \rightarrow \mathbb{RP}^2$. This map is almost surjective only missing the point that is the linear span of $(0, 0, 1)$ in \mathbb{RP}^2 . It is otherwise bijective and invertible exactly because any scalar multiple of a line is the same line and scalars leave you in the same linear span (point) in \mathbb{RP}^2 . This map shows that N is diffeomorphic to \mathbb{RP}^2 minus a point. Though I will not prove it here a little bit of algebraic topology on plane models of the projective plane and the mbius strip will yield that the projective plane minus a point is homeomorphic (and in fact diffeomorphic) to the mbius band without a boundary. \square

Note that the mbius band is not orientable but the cylinder is so N cannot be diffeomorphic to the cylinder.